



Guidelines for the Methodology of Cracked Tooth Studies

aae.org

Introduction

The guidelines for the methodology of cracked tooth epidemiologic studies are intended to allow institutions, practice-based research networks, large group practices and even individual private practitioners to collect and publish important data with regard to the incidence and/or prevalence of root cracks or fractures (RC/F) in teeth.

While they are not fixed protocols, the guidelines will standardize methodology and data collected across studies, facilitating future meta-analysis of the data from the studies that use this protocol. It should be noted that this methodology would not include cracks that result from acute dental trauma, such as a horizontal root fracture, but the increasingly common type that is perhaps a repetitive stress injury.



Special thanks to the Special Committee on
Methodology of Cracked Tooth Studies for
their work in developing these guidelines:

Shimon Friedman, D.M.D., Chair

Amir Azarpazhooh, D.D.S., M.Sc., Ph.D.

George A. Bruder III, D.M.D.

Keith V. Krell, D.D.S., M.S., M.A.

Isabel Mello, D.D.S., M.Sc., Ph.D.

Donald R. Nixdorf, D.D.S., M.S., Consultant

Robert S. Roda, D.D.S., M.S., Board Liaison

Methodology for Assessment of Prevalence of RC/F in Root-Filled Teeth

Eligible Study Designs

- Cross-sectional longitudinal study — follow [STROBE guidelines](#)

Methodology and Reporting Requirements

- Confirm approval of study protocol by relevant Institutional Review Board and compliance with informed-consent protocol for subject recruitment for the study.
- Estimate required sample size including reference data/assumptions.
- Define the study population of interest and describe methods of recruiting subjects.
- Specify eligibility criteria for subjects, if applied.
- Specify how root-filled teeth are identified, e.g., inspection of panoramic radiographic records, review of cone beam computed tomography volumes.
- Describe data collection process used to assess root-filled teeth, e.g., exposure of periapical radiographs, face-to-face interview, clinical examination, or combination of the above.
- Define inclusion/exclusion criteria for root-filled teeth, if applied. For example, to assess RC/F, time lapse of ≥ 2 years after endodontic treatment may be considered as threshold for inclusion.
- Define how missing teeth were considered, specifically how any root-filled teeth among them were identified.
- For root-filled missing teeth, describe how the study determined whether RC/F was the cause of extraction, e.g., by asking the patient, by examining treatment records, or by asking the dentist who last examined the tooth before extraction.
- Define outcome assessment/diagnostic (clinical, radiographic) measures of periapical health/disease and RC/F. Include specific features suggestive of/consistent with RC/F.*
- Specify outcome assessment criteria used, with specific mention of criteria for assessment as RC/F.*
- Define assessment (clinical, radiographic) measures and criteria for assessment of root-filling quality. Optional in study focused on RC/F.
- Define assessment (clinical, radiographic) measures and criteria for assessment of restoration type and quality. Optional in study focused on RC/F.
- Define interval period(s) between successive examinations of the same population. For assessment of RC/F, intervals of 5 to 10 years may be considered.

* See page 12 for list of specific diagnostic criteria for RC/F

Statistical Methods

- Define the approach to longitudinal data analysis and reporting, in regards to root-filled teeth captured at the inception of the study.
- Define method for univariate reporting of frequencies within the study sample.
- Define method for bivariate analysis of variables associated with the outcome(s) of interest, including prevalence of RC/F.
- Define method for multivariate analysis of outcome-associated variables.
- Define the level of significance.

Reporting of Results

- Report the study sample captured (N) at the outset of the study. Identify numbers of subjects, teeth, root-filled teeth, missing teeth.
- Characterize the study sample with regards to radiographic (and clinical, if assessed) findings.
- Report the numbers/frequencies of periapical health/disease and other variables of interest, i.e., root-filling quality, restoration type and quality.
- Report specifically on RC/F in captured teeth and, if construed, in missing teeth. Identify numbers/frequencies of the following:
 - teeth with obvious root fractures with separated fragments
 - teeth with fracture lines evident in radiographs or cone beam computed tomography images
 - teeth with radiographic findings suggestive of RC/F
 - teeth with clinical findings suggestive of RC/F
 - teeth with RC/F evident by direct inspection (observation of root surface, exploratory surgery, orthograde access, post-extraction)
- Report on the study sample captured (n) at each subsequent examination juncture, in regards to subjects, teeth, root-filled teeth, missing teeth, RC/F.
- Report specifically on changes observed within the subset of root-filled teeth, with regard to periapical health/disease, e.g., improvement, deterioration, no change, more missing teeth, RC/F.
- Where possible, in reporting of teeth diagnosed as having RC/F, differentiate between roots with and without posts.
- Report the bivariate analysis to identify variables associated with outcomes of interest, including RC/F.
- Report the multivariate analysis to identify predictive variables including those related to RC/F.

Methodology for Assessment of Incidence of RC/F in Root-Filled Teeth

Eligible Study Designs

- Prospective cohort study – follow [STROBE guidelines](#)
- Randomized controlled trial – follow [CONSORT guidelines](#)
- Retrospective cohort study – follow [STROBE guidelines](#)

Preoperative Data Collection and Reporting Requirements

- Confirm approval of study protocol by relevant Institutional Review Board and compliance with informed-consent protocol for subject recruitment for the study.
- Define inception cohort/study population/study groups.
- Define preoperative assessment/diagnostic (clinical and radiographic) measures and criteria.
- Specify inclusion/exclusion criteria, with specific mention of diagnostic features suggestive of/consistent with root crack/fracture.
- Define included study sample (N).
- Characterize the study sample in regards to demographic and pre-operative clinical and radiographic features.
- For randomized controlled trials, describe method of randomization for primary variable of interest and how secondary variables are controlled.
- Estimate required sample size including reference data/assumptions and projected attrition of the sample.

Intraoperative Data Collection and Reporting Requirements

- Describe all intervention steps/techniques/instruments/materials in detail, in a manner that will support duplication of the interventions by others. Include pertinent data regarding temporary and definitive restorations, including time elapsed between root filling and restoration.
- Describe intraoperative complications that occurred, if any.
- Outline the observation (follow-up) schedule and methods used to ascertain attendance, including incentives offered to subjects. The observation period(s) must be sufficient to express the outcome(s) of interest. For RC/F, this period could be 4-7 years or even longer.

Postoperative Data Collection and Reporting Requirements

- Define outcome assessment/diagnostic (clinical and radiographic) measures. Include specific features suggestive of/consistent with RC/F.*
- Differentiate RC/F from other types of tooth cracks and fractures (because the main dilemma about RC/F in root-filled teeth concerns roots that have no posts).
- Specify outcome assessment criteria, with specific mention of criteria for assessment as RC/F.*
- Describe methods used to characterize subjects lost-to-follow-up into categories of “dropouts” and “discontinuers.”
- Describe methods used to account for any teeth that have been lost or further treated (nonsurgically or surgically) during the observation period, including specific reasons that led to such occurrences.

* See page 12 for list of specific diagnostic criteria for RC/F.

Statistical Methods

- Define the approach(es) to data analysis and reporting, i.e., as one-point data, longitudinal data, incidence/frequency of health/disease or survival.
- Define method for univariate reporting of frequencies within the study cohort and sample.
- Define method for bivariate analysis of variables associated with the outcome(s) of interest.
- Define method for multivariate analysis of outcome-associated variables to identify outcome predictors.
- Define the level of significance.

Reporting of Results

- Define the final study sample (n) attending the end-point(s) of the study and characterize it in regards to variables of interest.
- Account for “dropouts” and “discontinuers” (whose absence is not assumed to be related to the interventions or outcomes of interest) and report the recall rate (%N).
- Characterize the final sample (n) in comparison to the original sample (N) and identify differences between the two samples, with regard to outcome predictors, to explore potential bias related to loss-to-follow-up.
- Report the number of teeth lost or further treated during the observation period and the reasons for these occurrences.
- Report the breakdown of results, including RC/F, in relation to specific outcome measures or the outcome criteria or both. Report specifically on teeth diagnosed as having RC/F while differentiating between roots with and without posts.
- Report the bivariate analysis to identify potential outcome predictors, including potential predictors of RC/F.
- Report the multivariate analysis to identify outcome predictors, including predictors of RC/F.

Template for Data Collection

Type of Data Possible Entries

Demographic Data	
Sex	<input type="checkbox"/> Female <input type="checkbox"/> Male
Age (years)	<input type="checkbox"/> 15-24 <input type="checkbox"/> 25-34 <input type="checkbox"/> 35-44 <input type="checkbox"/> 45-54 <input type="checkbox"/> 55-64 <input type="checkbox"/> ≥65
Treated tooth	(enter number 1-32) _____
Preoperative Clinical Symptoms and Signs	
Spontaneous pain	<input type="checkbox"/> Absent <input type="checkbox"/> Present
Triggered pain	<input type="checkbox"/> Biting <input type="checkbox"/> Touch <input type="checkbox"/> Cold <input type="checkbox"/> Hot <input type="checkbox"/> Sweet
Swelling	<input type="checkbox"/> Absent <input type="checkbox"/> Buccal <input type="checkbox"/> Lingual/palatal
Sinus tract	<input type="checkbox"/> Absent <input type="checkbox"/> Buccal <input type="checkbox"/> Lingual/palatal
Preoperative Diagnostic Data – Clinical	
Cold test	<input type="checkbox"/> Positive <input type="checkbox"/> Non-lingering <input type="checkbox"/> Lingering <input type="checkbox"/> Negative
Heat test	<input type="checkbox"/> No pain elicited <input type="checkbox"/> Pain elicited
Percussion	<input type="checkbox"/> Not tender <input type="checkbox"/> Tender <input type="checkbox"/> Very tender
Palpation	<input type="checkbox"/> Not tender <input type="checkbox"/> Tender
Mobility	<input type="checkbox"/> Physiological <input type="checkbox"/> 1 <input type="checkbox"/> 2 <input type="checkbox"/> 3
Probing depth	<input type="checkbox"/> ≤ 3 mm <input type="checkbox"/> 4-5 mm <input type="checkbox"/> ≥ 6 mm
Probed defect location	<input type="checkbox"/> Mesial <input type="checkbox"/> Distal <input type="checkbox"/> Buccal <input type="checkbox"/> Lingual <input type="checkbox"/> None
Tooth Slooth	<input type="checkbox"/> No pain <input type="checkbox"/> Pain at one cusp <input type="checkbox"/> Pain at ≥2 cusps
Coronal crack	<input type="checkbox"/> Not evident <input type="checkbox"/> Buccal <input type="checkbox"/> Lingual/palatal
Root crack (with gingiva reflected)	<input type="checkbox"/> Not evident <input type="checkbox"/> Buccal <input type="checkbox"/> Lingual/palatal
Fractured/dis-lodged restoration	<input type="checkbox"/> Not evident <input type="checkbox"/> Evident
Preoperative Radiographic Findings	
Periapical area of radiolucency (low attenuation)	<input type="checkbox"/> Absent <input type="checkbox"/> Widened PDL space <input type="checkbox"/> 2-4 mm (widest dimension) <input type="checkbox"/> 5-7 mm (widest dimension) <input type="checkbox"/> ≥ 8 mm (widest dimension)
Lateral area of radiolucency (enter applicable roots)	<input type="checkbox"/> Absent <input type="checkbox"/> Widened PDL space <input type="checkbox"/> Apical 1/3 <input type="checkbox"/> Middle 1/3 <input type="checkbox"/> Coronal 1/3 <input type="checkbox"/> Entire root length
Furcal area of radiolucency	<input type="checkbox"/> Absent <input type="checkbox"/> Level of coronal 1/3 <input type="checkbox"/> Level of middle 1/3 <input type="checkbox"/> Level of apical 1/3 <input type="checkbox"/> Entire root length
Root fracture	<input type="checkbox"/> Not evident <input type="checkbox"/> Evident

Type of Data Possible Entries

Preoperative Diagnosis						
Pulp	<input type="checkbox"/> Normal	<input type="checkbox"/> Reversible pulpitis	<input type="checkbox"/> Asymptomatic irreversible pulpitis	<input type="checkbox"/> Symptomatic irreversible pulpitis	<input type="checkbox"/> Necrosis	<input type="checkbox"/> Previously treated
Apical	<input type="checkbox"/> Normal	<input type="checkbox"/> Asymptomatic apical periodontitis	<input type="checkbox"/> Symptomatic apical periodontitis	<input type="checkbox"/> Chronic apical abscess	<input type="checkbox"/> Acute apical abscess	
Root	<input type="checkbox"/> Intact	<input type="checkbox"/> Vertical crack suspected	<input type="checkbox"/> Vertical crack/fracture	<input type="checkbox"/> Horizontal fracture suspected	<input type="checkbox"/> Horizontal fracture	
Intraoperative (intervention) Data						
Preflaring	<input type="checkbox"/> Gates-Glidden drills	<input type="checkbox"/> Orifice Shapers	<input type="checkbox"/> Other: _____		<input type="checkbox"/> None	
Instrumentation	<input type="checkbox"/> Hand instruments only	<input type="checkbox"/> Rotary used	<input type="checkbox"/> Reciprocation used	<input type="checkbox"/> Other (specify) _____		
Irrigation (check all that apply)	NaOCl <input type="checkbox"/> 1% <input type="checkbox"/> 2.5% <input type="checkbox"/> 5%	EDTA <input type="checkbox"/> 17% <input type="checkbox"/> Other ____ %	Chlorhexidine <input type="checkbox"/> 0.12% <input type="checkbox"/> 2%	<input type="checkbox"/> MTAD	<input type="checkbox"/> QMix	<input type="checkbox"/> Other (specify) _____
Intracanal medication	<input type="checkbox"/> Calcium hydroxide	<input type="checkbox"/> Other (specify) _____			<input type="checkbox"/> None	
Medication period	<input type="checkbox"/> < 7 days	<input type="checkbox"/> 7-10 days	<input type="checkbox"/> 11-14 days	<input type="checkbox"/> > 14 days	<input type="checkbox"/> None	
MAF sizes (enter for each canal)	<input type="checkbox"/> Distal/palatal	<input type="checkbox"/> Mesio-buccal/buccal	<input type="checkbox"/> Mesio-lingual/lingual/MB2	<input type="checkbox"/> Disto-buccal	<input type="checkbox"/> Single	<input type="checkbox"/> Other (specify) _____
Root filling technique	<input type="checkbox"/> Cold lateral	<input type="checkbox"/> Warm lateral	<input type="checkbox"/> Warm vertical	<input type="checkbox"/> Carrier based	<input type="checkbox"/> Single cone	<input type="checkbox"/> Other (specify) _____
Temporary access restoration	<input type="checkbox"/> Composite resin	<input type="checkbox"/> Glass-ionomer cement	<input type="checkbox"/> IRM	<input type="checkbox"/> Cavit	Cotton pellet placed: <input type="checkbox"/> Yes <input type="checkbox"/> No	
Final restoration	<input type="checkbox"/> Glass-ionomer cement	<input type="checkbox"/> Composite resin	<input type="checkbox"/> Amalgam	<input type="checkbox"/> Onlay	<input type="checkbox"/> Crown	
Timing of final restoration	<input type="checkbox"/> Immediate	<input type="checkbox"/> ≤ 2 weeks	<input type="checkbox"/> ≤ 2 - 4 weeks	<input type="checkbox"/> > 4 weeks		

(Continued on next page)

Template for Data Collection

Type of Data Possible Entries

Intraoperative (intervention) Data (continued)						
Post	<input type="checkbox"/> Absent	<input type="checkbox"/> Cast	<input type="checkbox"/> Prefabricated metallic	<input type="checkbox"/> Prefabricated fiber	<input type="checkbox"/> Prefabricated ceramic	
Post extent (relative to crestal bone)	<input type="checkbox"/> 1-2 mm	<input type="checkbox"/> 3-4 mm	<input type="checkbox"/> 5-6 mm	<input type="checkbox"/> > 6 mm		
Post width	<input type="checkbox"/> ≤ 1/3 of root width	<input type="checkbox"/> 1/2 of root width	<input type="checkbox"/> ≥ 3/4 of root width			
Post luting cement	<input type="checkbox"/> Dentin-bonded		<input type="checkbox"/> Non-bonded			
Procedural complication	Perforation: <input type="checkbox"/> chamber <input type="checkbox"/> coronal 1/3 <input type="checkbox"/> middle 1/3 <input type="checkbox"/> apical 1/3		Instrument fracture: <input type="checkbox"/> coronal 1/3 <input type="checkbox"/> middle 1/3 <input type="checkbox"/> apical 1/3		Crack extending into canal: <input type="checkbox"/> distal/palatal <input type="checkbox"/> mesio-buccal/buccal <input type="checkbox"/> mesio-lingual/lingual <input type="checkbox"/> disto-buccal <input type="checkbox"/> other	
Postoperative (follow-up) Clinical Diagnostic Data						
Observation period	<input type="checkbox"/> < 1 year	<input type="checkbox"/> 1-2 years	<input type="checkbox"/> > 2-3 years	<input type="checkbox"/> > 3-4 years	<input type="checkbox"/> > 4-5 years	<input type="checkbox"/> > 5 years
Further treatment	<input type="checkbox"/> Nonsurgical	<input type="checkbox"/> Apical surgery	<input type="checkbox"/> Root amputation	<input type="checkbox"/> Hemisection	<input type="checkbox"/> Intentional replantation	<input type="checkbox"/> Extraction
Spontaneous pain	<input type="checkbox"/> Absent		<input type="checkbox"/> Present			
Triggered pain on biting	<input type="checkbox"/> Absent		<input type="checkbox"/> Present			
Swelling	<input type="checkbox"/> Absent		<input type="checkbox"/> Present			
Sinus tract	<input type="checkbox"/> Absent	<input type="checkbox"/> Buccal	<input type="checkbox"/> Lingual/palatal			
Percussion	<input type="checkbox"/> Negative		<input type="checkbox"/> Positive			
Palpation	<input type="checkbox"/> Negative		<input type="checkbox"/> Positive			
Mobility	<input type="checkbox"/> Physiological	<input type="checkbox"/> 1	<input type="checkbox"/> 2	<input type="checkbox"/> 3		
Probing depth	<input type="checkbox"/> ≤ 3 mm	<input type="checkbox"/> 4-5 mm	<input type="checkbox"/> ≥ 6 mm			
Probed defect location	<input type="checkbox"/> Mesial	<input type="checkbox"/> Distal	<input type="checkbox"/> Buccal	<input type="checkbox"/> Lingual	<input type="checkbox"/> None	
Root crack (with gingiva reflected)	<input type="checkbox"/> Not evident		<input type="checkbox"/> Buccal	<input type="checkbox"/> Lingual/palatal		
Fractured/dislodged restoration	<input type="checkbox"/> Not evident		<input type="checkbox"/> Evident			

Type of Data Possible Entries

Postoperative Radiographic Findings						
Periapical area of radiolucency (low attenuation)	<input type="checkbox"/> Absent	<input type="checkbox"/> Widened PDL space	<input type="checkbox"/> 2-4 mm (widest dimension)	<input type="checkbox"/> 5-7 mm (widest dimension)	<input type="checkbox"/> ≥ 8 mm (widest dimension)	
Lateral area of radiolucency (enter applicable roots)	<input type="checkbox"/> Absent	<input type="checkbox"/> Widened PDL space	<input type="checkbox"/> Apical 1/3	<input type="checkbox"/> Middle 1/3	<input type="checkbox"/> Coronal 1/3	<input type="checkbox"/> Entire root length
Furcal area of radiolucency	<input type="checkbox"/> Absent	<input type="checkbox"/> Level of coronal 1/3	<input type="checkbox"/> Level of middle 1/3	<input type="checkbox"/> Level of apical 1/3	<input type="checkbox"/> Entire root length	
Root fracture	<input type="checkbox"/> Not evident	<input type="checkbox"/> Evident				
Postoperative CBCT Findings						
Root fracture/separation	<input type="checkbox"/> Not evident	<input type="checkbox"/> Mesial	<input type="checkbox"/> Distal	<input type="checkbox"/> Buccal	<input type="checkbox"/> Lingual/palatal	
Bone defect pattern	<input type="checkbox"/> Lateral - narrow	<input type="checkbox"/> Partial root length	<input type="checkbox"/> Total root length	<input type="checkbox"/> Bone plate eroded		
Postoperative Diagnosis – Treatment Outcome						
Apical	<input type="checkbox"/> Normal	<input type="checkbox"/> Asymptomatic apical periodontitis	<input type="checkbox"/> Symptomatic apical periodontitis	<input type="checkbox"/> Chronic apical abscess	<input type="checkbox"/> Acute apical abscess	
Root	<input type="checkbox"/> Intact	<input type="checkbox"/> Vertical crack suspected	<input type="checkbox"/> Vertical crack/fracture	<input type="checkbox"/> Horizontal fracture suspected	<input type="checkbox"/> Horizontal fracture	
Cracked/fractured root	<input type="checkbox"/> Without post	<input type="checkbox"/> With post	<input type="checkbox"/> Mesial/mesio-buccal	<input type="checkbox"/> Distal/disto-buccal	<input type="checkbox"/> Palatal/lingual	<input type="checkbox"/> Buccal <input type="checkbox"/> Single

Diagnostic Criteria for Application in Epidemiological Studies on RC/F in Root-Filled Teeth

Listed features may be used to diagnose or differentially diagnose RC/F.

Diagnosed as RC/F	Differentially Diagnosed as RC/F	Comments
Observed Features – Clinical		
Spontaneous pain	Spontaneous pain	
Pain on biting	Pain on biting	
Swelling	Swelling	
Single sinus tract	Single sinus tract	
Buccal + lingual/palatal sinus tracts*		
Percussion tenderness	Percussion tenderness	
Palpation tenderness	Palpation tenderness	
Increased mobility	Increased mobility	Mobility 2 or 3
Narrow isolated probing ≥ 6 mm	Narrow isolated probing ≥ 6 mm	Without periodontal disease
Buccal + lingual narrow probing ≥ 6 mm*		Without periodontal disease
Root crack evident*		With gingiva reflected, staining, transillumination, magnification
Observed Features – Radiographic		
Root fracture/separation evident*		
"J" shape defect	"J" shape defect	Without periodontal disease
Extensive radiolucency	Extensive radiolucency	≥ 5 mm
Lateral radiolucency	Lateral radiolucency	<ul style="list-style-type: none"> • Apical 1/3, middle 1/3, coronal 1/3, entire root length • Without periodontal disease
Lateral widened PDL space*		
Furcal radiolucency	Furcal radiolucency	<ul style="list-style-type: none"> • Extends to middle 1/3 or entire root length • Without periodontal disease

* Typical feature of RC/F

Diagnosed as RC/F

Differentially Diagnosed as RC/F

Comments

Observed Features – Limited Field of View CBCT		
Root fracture/separation evident*		
Single lateral narrow radiolucency	Lateral narrow radiolucency	<ul style="list-style-type: none"> • Apical 1/3, middle 1/3, coronal 1/3, entire root length • Without periodontal disease
Buccal + lingual lateral narrow radiolucency*		<ul style="list-style-type: none"> • Apical 1/3, middle 1/3, coronal 1/3, entire root length • Without periodontal disease
Furcal radiolucency	Furcal radiolucency	Without periodontal disease
Loss of cortical plate	Loss of cortical plate	Full length of root
Radiolucency surrounding entire root		
Observed Features – Exploratory		
Crack line detected upon exploration*		Extraction, surgical exposure or endodontic access



Email: info@aae.org

f facebook.com/endodontists

t [@SavingYourTeeth](https://twitter.com/SavingYourTeeth)

YouTube youtube.com/rootcanalspecialists

W www.aae.org

© 2016

American Association of Endodontists (AAE),
All Rights Reserved